Modern Physics Letters A
© World Scientific Publishing Company

arXiv:1612.01269v3 [gr-qc] 16 Jan 2017

On the Time Dependent Rindler Hamiltonian Single Particle Eigen States in Momentum Space

Soma Mitra

[a] *Department of Physics, Visva-Bharati, Santiniketan, India 731235*
E-mail: somaphysics@gmail.com

Sanchita Das

[a] *Department of Physics, Visva-Bharati, Santiniketan, India 731235*
E-mail: sanchedasm@gmail.com

Somenath Chakrabarty[a]

[a] *Department of Physics, Visva-Bharati, Santiniketan, India 731235*
E-mail: somenath.chakrabarty@visva-bharati.ac.in

Received (Day Month Year)
Revised (Day Month Year)

We have developed a formalism to get the time evolution of the eigen states of Rindler Hamiltonian in momentum space. We have discussed the difficulties with characteristic curves, and re-cast the time evolution equations in the form of two-dimensional Laplace equation. The solutions are obtain both in polar coordinates as well as in the Cartesian form. It has been observed that in the Cartesian coordinate, the probability density is zero both at $t = 0$ (the initial time) and at $t = \infty$ (the final time) for a give x-coordinate. The reason behind such peculiar behavior of eigen state is because it satisfies 1+1-dimensional Laplace equation.

Keywords: Uniformly accelerated motion; Rindler coordinates; Principle of equivalence; Laplace equation; Schriödinger equation in momentum space

PACS Nos.: 03.65.Ge,03.65.Pm,03.30.+p,04.20.-q

Exactly like the flat Minkowski space-time geometry, the Rindler space is also locally flat.[1–8] The only difference is that the observer is moving with uniform acceleration.[1–8] Then according to the principle of equivalence, the Rindler space corresponds to a frame at rest in presence of a locally uniform gravitational field. The strength of the gravitational field is exactly equal to the magnitude of the local acceleration.[1,2,9,10] The space-time transformations in Rindler space are given by

$$ct = \left(\frac{c^2}{\alpha} + x'\right) \sinh\left(\frac{\alpha t'}{c}\right) \quad \text{and}$$
$$x = \left(\frac{c^2}{\alpha} + x'\right) \cosh\left(\frac{\alpha t'}{c}\right)$$

Hence the Rindler metric can be written as

$$g^{\mu\nu} \equiv \text{diag}\left(\left(1+\frac{\alpha x}{c^2}\right), -1, -1, -1\right)$$

In $1+1$ dimension it reduces to

$$g^{\mu\nu} \equiv \text{diag}\left(\left(1+\frac{\alpha x}{c^2}\right), -1\right)$$

Then with some simple algebra it can be shown that the single particle Hamiltonian in Rindler space in one-dimensional form is given by[5–8]

$$H = \left(1+\frac{\alpha x}{c^2}\right)(p^2c^2 + m_0^2c^4)^{1/2} \tag{1}$$

where α is the uniform acceleration of the frame or the uniform gravitational field in the neighborhood of the point indicated by the coordinate x. Here p and m_0 are the particle momentum along the positive x-direction and the rest mass respectively and c is the speed of light in free space.

Further, the observer is assumed to have one dimensional motion along positive x-direction. Then in the non-relativistic approximation, neglecting the rest mass energy term, the Hamiltonian given by eqn.(1) reduces to

$$H = \left(1+\frac{\alpha x}{c^2}\right)\frac{p^2}{2m_0} \tag{2}$$

Now in the quantum mechanical scenario, the classical dynamical variables x, p and H are treated as operators and the canonical quantization condition $[x,p] = i\hbar$ is satisfied. The aim of this article is to study the evolution of Rindler Hamiltonian eigenstate with time in momentum space in $1+1$ dimension. This is the continuation of some of our previously reported results.[6,7,11] Then following the article by R.W. Robinett in Am. Jour. Phys.,[12] the time dependent Schrödinger equation with the momentum representation of the positional coordinate operator x ($x = i\hbar\partial/\partial p$) is given by

$$\left(1+\frac{\alpha}{c^2}i\hbar\frac{\partial}{\partial p}\right)\frac{p^2}{2m_0}\psi(p,t) = i\hbar\frac{\partial\psi(p,t)}{\partial t} \tag{3}$$

After the substitution of

$$\frac{p^2}{2m_0}\psi(p,t) = \phi(p,t) \tag{4}$$

it becomes

$$\phi(p.t) + i\hbar\frac{\alpha}{c^2}\frac{\partial\phi(p,t)}{\partial p} = i\hbar\frac{2m_0}{p^2}\frac{\partial\phi(p,t)}{\partial t} \tag{5}$$

To solve this equation analytically, we multiply both sides of the above differential equation by the integrating factor

$$I_p = \exp\left(\frac{i\hbar\alpha}{c^2}p\right) \tag{6}$$

Then the above Schrödinger equation reduces to

$$\frac{\partial \xi}{\partial \tau} - i\frac{\partial \xi}{\partial q} = 0 \tag{7}$$

where

$$\xi(q,\tau) = I_p\phi(p,t), \quad \tau = \frac{t}{2m_0\hbar} \text{ and } q = \frac{1}{p} \tag{8}$$

Let us now try to solve this reduced form of the Schrödinger equation given by eqn.(7). We first try with the well known characteristics flow equation approach.[13] We introduce two new variables

$$r = \tau + iq \text{ and } s = -i\tau - q \tag{9}$$

Then the above differential equation reduces to

$$\frac{\partial \xi(r,s)}{\partial s} = 0 \tag{10}$$

which gives $\xi(\tau, q) = f(r) = f(\tau + iq)$ Here $f(\tau + iq)$ is completely arbitrary. The arbitrariness of the solution can sometime be removed by the boundary conditions. However, the complex characteristics given by eqn.(10) has no practical value. Also it has no obvious relevance to boundary conditions. Moreover, the propagation along such complex characteristics will not provide any acceptable partial differential equation for a real physical system. Or in other wards there will be no acceptable form of differential equation to represent the flow with physical boundary conditions. Therefore we shall try an alternative approach to get the solution.

We follow the technique to obtain Cauchy-Euler differential equation.[13,14] We take derivatives of both the terms of eqn.(7) with respect to τ and $-iq$ and then subtracting these two equations, we get

$$\frac{\partial^2 \xi}{\partial \tau^2} + \frac{\partial^2 \xi}{\partial q^2} = 0 \tag{11}$$

This is the well known Laplace equation in two-dimension, also known as the elliptic form of partial differential equation. However, for $a = b$, we may call the above equation as circular type partial differential equation. To get solutions of this equation we consider circular polar coordinate (ρ, ϕ), where $\tau = \rho\cos\phi$, $q = \rho\sin\phi$ and $\phi = \tan^{-1}(q/\tau) = \tan^{-1}(\text{constant}/pt)$. Then the above differential equation reduces to

$$\frac{1}{\rho}\frac{\partial}{\partial \rho}\left(\rho\frac{\partial \xi}{\partial \rho}\right) + \frac{1}{\rho^2}\frac{\partial^2 \xi}{\partial \phi^2} = 0 \tag{12}$$

With the separable form of wave function, given by $\xi(\rho, \phi) = R(\rho)\Phi(\phi)$, we have

$$\frac{d^2\Phi}{d\phi^2} + \nu^2\Phi = 0 \tag{13}$$

and

$$\rho\frac{d}{d\rho}\left(\frac{dR}{d\rho}\right) - \nu^2 R = 0 \tag{14}$$

where ν is an unknown real constant. Here the first one is the well known differential equation for classical harmonic oscillator in one dimension, whereas the second one is called the Cauchy-Euler equation. The solution of the first equation is

$$\Phi(\phi) = A\cos(\nu\phi) + B\sin(\nu\phi) \tag{15}$$

with A and B are two unknown constants. The solutions of the Cauchy-Euler equation are known to be of the form ρ^{γ}. Substituting in eqn.(15), we have $\gamma = \pm\nu$. Hence for the radial part we may write

$$R(\rho) = a\rho^{\nu} + b\rho^{-\nu} \tag{16}$$

where a and b are two other unknown constants. In fig.(1) we have plotted for the sake of illustration the wave function $\xi(\rho, \phi)$ for all the constants equal to unity along with $\nu = 1$. The variable ϕ is along x-axis and ρ is along $y - axis$. One should note that from the definitions of ϕ and ρ it is quite obvious that both of them are positive in nature.



Fig. 1. Variation of $\xi(\rho, \phi)$ with ϕ, along x-axis and ρ along y-axis.

As a special case, for $\nu = 0$

$$\Phi(\phi) = A_0 + B_0\phi \tag{17}$$

and

$$R(\rho) = a_0 + b_0 \ln \rho \tag{18}$$

In fig.(2) we have plotted the wave function $\xi(\rho, \phi)$ for this special case. In the electrostatic problem the above solutions are generally considered as the building blocks to construct the potential by linear superposition. The radial logarithmic form gives the potential due to a linear charge distribution. Taking into account this concept of superposition principle, we can also write down the general solution for the wave functions. However, in the present situation, ϕ is restricted in the first

quadrant only ($0 \leq \phi \leq \pi/2$). Therefore, ν does not necessarily have integer values. Therefore no further progress can be achieved from this approach.

Next we shall try to get solutions in $\tau - q$ space, which is equivalent to two dimensional $(x-y)$ Cartesian coordinate system. Using the separable form of solutions $\xi(\tau, q) = T(\tau)Q(q)$, we have

$$\frac{d^2T}{d\tau^2} + \omega^2 T = 0 \tag{19}$$

and

$$\frac{d^2Q}{dq^2} - \omega^2 Q = 0 \tag{20}$$

where ω is an unknown real constant. It is trivial to show that the solutions are

$$T(\tau) = C_1 \sin\omega\tau + C_2 \cos\omega\tau \tag{21}$$

and

$$Q(q) = D_1 \sinh\omega q + D_2 \cosh\omega q \tag{22}$$

Where C_1, C_2, D_1 and D_2 are unknown constants. Since τ is proportional to the actual time coordinate t, which increases continuously, and not periodic or bounded, there is no problem in having same wave function at various time coordinate. Therefore, ω is not necessarily an integer. Whereas $q \propto 1/p$, therefore for large p, $q \longrightarrow 0$ and then neglecting the higher degree terms, $Q \propto q \propto 1/p$. In the case of $p \longrightarrow 0$, i.e., $q \longrightarrow \infty$, the asymptotic form of the solution $Q \longrightarrow \infty$, which is unphysical. To make the solution physically acceptable, we put by hand $D_2 = -D_1 = -D$, then

$$Q(q) = D(\cosh\omega q - \sinh\omega q) \tag{23}$$

Which becomes after re-defining the constant D,

$$Q(q) = D\exp(-\omega q) \tag{24}$$

This is the asymptotic form of the solution. Therefore the final solutions for all possible values of ω are given by

$$\xi(\tau, q) = \sum_{\omega}(C_1 \sin\omega\tau + C_2 \cos\omega\tau)(D_1 \sinh\omega q + D_2 \cosh\omega q) \tag{25a}$$

and asymptotically

$$\xi(\tau, q) = \sum_{\omega}(C_1 \sin\omega\tau + C_2 \cos\omega\tau)\exp(-\omega q) \tag{25b}$$

Assuming that the variable ω is changing continuously, we can transform the sum over ω by integral, and then we can write down the asymptotic form of the wave function after evaluating the integral over ω within the limit 0 to ∞ and is given by:

$$\xi(\tau, q) = \frac{C_1\tau + C_2 q}{\tau^2 + q^2} \tag{25}$$

where C_1 and C_2 are two re-defined constants to make the expression dimensionally correct. In fig.(3) we have plotted $\xi(\tau, q)$ for $\omega = 1$ and all the constants are also set equal to unity. In fig.(4) we have shown the same wave function, but in the asymptotic region. Whereas in fig.(5) we have plotted the asymptotic form of wave function for all possible values of ω, which is assumed to be changing continuously.

Therefore, it is quite obvious that the time dependence of eigen state of the Rindler Hamiltonian is quite different from the stationary state solutions of the wave functions. To make it more clear we consider the Fourier transform of eqn.(25) to obtain the space-time dependent form of the wave function. The Fourier transform is given by

$$\psi(x,t) = \frac{N}{2\pi}\int_{-infty}^{+\infty} \frac{at + \frac{b}{p}}{a^2t^2 + \frac{b^2}{p^2}} \exp(-ipx)dp \tag{26}$$

where N is the normalization constant. On evaluating the Fourier transform, we get

$$\psi(x,t) = N\frac{b}{2^{1/2}a^2t^2} \exp\left(-\frac{bx}{at} - i\frac{\pi}{4}\right) \tag{27}$$

The normalized form of the wave function is given by

$$\psi(x,t) = \left(\frac{2b}{at}\right)^{1/2} \exp\left(-\frac{bx}{at} - i\frac{\pi}{4}\right) \tag{28}$$

The probability density is given by

$$\mid \psi(x,t) \mid^{1/2} = \frac{2b}{at} \exp\left(-\frac{2bx}{at}\right) \tag{29}$$

Obviously the probability density, i.e., the probability of existence of the particle becomes zero for both $t \longrightarrow 0$ and $t \longrightarrow \infty$. Then in between there is some time at which the probability of existence becomes maximum, given by

$$\frac{d}{dt} \mid \psi \mid^2 = 0 \tag{30}$$

This condition gives

$$t = \frac{2bx}{a} = t_{max} \tag{31}$$

This is the time at which the probability of existence becomes maximum. In fig.(6) we have plotted the scaled quantity

$$\frac{\mid \psi(x,t) \mid}{(2b/a)^{1/2}} = \frac{\exp(-\gamma/t)}{t^{1/2}} \tag{32}$$

where $\gamma = bx/a$. In fig.(6) top one curve is for $\gamma = 0.1$, rest are in sequence $\gamma = 0.5$, 1, 5, 10 and 25 respectively. The type time dependence as shown above is because the wave functions satisfy the Laplace equation (eqn.(11)) in two-dimension. Finally we conclude by saying that the time dependencies of the Rindler Hamiltonian eigen state is quite different from the usual stationary state wave functions.

References

1. N.D. Birrell and P.C.W. Davies, Quantum Field Theory in Curved Space, Cambridge University Press, Cambridge, (1982).
2. W. Rindler, Essential Relativity, Springer-Verlag, New York, (1977).
3. G. F. Torres del Castillo and C.L. Perez Sanchez, Revista Mexican De Fisika **52**, 70, (2006).
4. Domingo J. Louis-Martinez, arXiv:10123063.
5. M. Socolovsky, Annales de la Foundation Louis de Broglie **39**, 1, (2014).
6. Sanchari De, Sutapa Ghosh and Somenath Chakrabarty, Modern Physics Letter A, **30**, 1550182, (2015).
7. Sanchari De, Sutapa Ghosh and Somenath Chakrabarty, Astrophys. Space Sci., (2015) 360:8, DOI 10.1007/s10509-015-2520-3.
8. C-G Huang, and J-R, Sun, J-R, arXiv:gr-qc/0701078, (2007).
9. Landau L.D. and Lifshitz E.M., The Classical Theory of Fields, Butterworth-Heimenann, Oxford, (1975).
10. C.W. Misner, Kip S. Thorne and J.A. Wheeler, Gravitation, W.H. freeman and Company, New York, (1972).
11. Soma Mitra and Somenath Chakrabarty, EPJ Plus (Communicated 2017).
12. R.W. Robinett, Am. Jour. Phys., **64**, 803, (1996).
13. Mathematical Methods for Physicists, G.B. Arfken, H.J. Weber and F.E. Harris, Seventh Editon, Elsevier, New York, (2013).
14. Mathematical Physics- A.Modern Introduction to Its Foundation, Sadri Hassani, Springer, New York, (1998).



Fig. 2. The surface plot of the wave function. ϕ is along x-axis and ρ along y-axis.



Fig. 3. The surface plot of the wave function $\xi(\tau, q)$, tau is along x-direction and q along y-axis.



Fig. 4. The surface plot of the asymptotic form of the wave function $\xi(\tau, q)$, tau is along x-direction and q along y-axis.



Fig. 5. The surface plot of the asymptotic form of the wave function $\xi(\tau, q)$, *tau* is along x-direction and q along y-axis.



Fig. 6. The variation of the scaled $\mid \psi(x,t) \mid$ (along y-axis) with time t (along x-axis) for various values of γ as mentioned in the text.